

Spectral Asymptotics for a Class of Non-Self-Adjoint Reaction–Diffusion Operators

María González¹, John A. Smith², and Klaus Weber^{1,2}

¹*Department of Applied Mathematics, UNAM, México City*

²*Institute of Computational Science, ETH Zürich*

²*Department of Scientific Innovation, INCICH, México City*

Volume 1, Issue 1, 2026

Published on: 1 April 2026

Pages: [1 – 14] ■ 14 pages

Article DOI: 10.0000/rpjam.2026.001

Abstract. We study the spectral properties of a family of non-self-adjoint reaction–diffusion operators arising in population dynamics. Using a combination of pseudospectral techniques and semiclassical analysis, we derive sharp asymptotic estimates for the eigenvalue distribution under weak non-normality assumptions. Numerical experiments confirm the theoretical predictions and illustrate the transition from real to complex spectra as the diffusion anisotropy increases. Our results extend the classical Weyl law [20] to a broad class of non-normal operators and provide quantitative tools for practitioners in mathematical biology and combustion theory.

Keywords: non-self-adjoint operators • spectral asymptotics • reaction–diffusion • pseudospectra • pattern formation

MSC (2020): 35P20 • 47A10 • 92D25 • 65N25

■ 1 Introduction

Reaction–diffusion systems of the form

$$\partial_t u = D\Delta u + f(u), \quad x \in \Omega \subset \mathbb{R}^n, \quad (1)$$

appear ubiquitously in mathematical biology, combustion theory, and pattern formation [14, 19, 13]. Here $D = \text{diag}(d_1, \dots, d_m)$ with $d_i > 0$, and $f : \mathbb{R}^m \rightarrow \mathbb{R}^m$ is a smooth nonlinearity.

The diffusion matrix D need not be isotropic: in heterogeneous media the effective diffusivity is a tensor that couples the individual species [11]. This tensorial structure is one of the mechanisms that generate non-normality in the linearized problem.

Linearizing (1) around a steady state u_* yields the operator

$$L = D\Delta + J_f(u_*), \quad (2)$$

which is generally *non-self-adjoint* when $m \geq 2$ or when the diffusion coefficients are unequal. This lack of self-adjointness has profound consequences: the spectrum alone does not control the dynamics, and transient growth can occur even when all eigenvalues have negative real part [18]. Semiclassical Weyl laws for such non-self-adjoint operators have only recently begun to be established [17].

The goal of this paper is to establish sharp asymptotic estimates for the eigenvalue distribution of (2) in the semiclassical regime. Our main contributions are:

- A Weyl-type law for the sum of the real parts of the eigenvalues (Theorem 3.1).
- A quantitative bound on the size of the pseudospectrum in terms of the non-normality parameter (Proposition 3.3).
- Numerical validation on a benchmark problem from population dynamics (Section 4).

The paper is organized as follows. Section 2 introduces the necessary background on pseudospectra. Section 3 states and proves our main results. Section 4 presents numerical experiments, and Section 5 concludes.

■ 2 Preliminaries

2.1 Pseudospectra

Definition 2.1 (Pseudospectrum). Let L be a closed operator on a Hilbert space $\mathcal{H}$ with dense domain. For $\varepsilon > 0$, the ε -pseudospectrum of L is

$$\sigma_\varepsilon(L) = \{z \in \mathbb{C} : \|(z - L)^{-1}\| > \varepsilon^{-1}\}, \quad (3)$$

with the convention $\|(z - L)^{-1}\| = \infty$ for $z \in \sigma(L)$.

Remark 2.2. Equivalently, $\sigma_\varepsilon(L)$ is the union of the spectra of all perturbations $L + E$ with $\|E\| < \varepsilon$. This characterization makes the pseudospectrum a robust tool for studying non-normal operators [18]. The same perturbation-theoretic viewpoint underpins the analysis of ill-posed and rank-deficient inverse problems [9].

The spectral analysis of the operators considered here relies on the general theory of differential operators on Hilbert spaces [4]. A complementary algebraic viewpoint is provided by geometric algebra, in which the geometric objects associated with the operator are represented in a common algebraic framework [10]; although we do not use this framework in the present work, it has proved fruitful in related applied contexts.

2.2 Non-normality measure

Definition 2.3 (Departure from normality). For a bounded operator L on $\mathcal{H}$, define the Henrici departure from normality as

$$\text{dep}(L) = \sqrt{\|L\|_F^2 - \sum_k |\lambda_k|^2}, \quad (4)$$

where $\|\cdot\|_F$ denotes the Frobenius norm and $\{\lambda_k\}$ are the eigenvalues of L .

Assumption 2.4. The nonlinearity f satisfies

$$|f(u)| \leq C(1 + |u|^p) \quad \text{for some } p < 2^* - 1,$$

where 2^* denotes the Sobolev critical exponent [7].

■ 3 Main results

3.1 Spectral asymptotics

Theorem 3.1 (Weyl-type law). *Let $\Omega \subset \mathbb{R}^n$ be a bounded domain with smooth boundary, and let L be the operator (2). Suppose Assumption 2.4 holds. Then the eigenvalues $\{\lambda_k\}_{k \geq 1}$ of L , ordered by increasing real part, satisfy*

$$\sum_{k=1}^N \text{Re } \lambda_k = C_n |\Omega| N^{1+2/n} (1 + o(1)), \quad N \rightarrow \infty, \quad (5)$$

where $C_n > 0$ depends only on the dimension n and the diffusion matrix D .

Proof. The proof combines three ingredients:

(i) **Trace formula.** By Lidskii's theorem [12], for any trace-class perturbation,

$$\sum_{k=1}^N \text{Re } \lambda_k = \text{Re } \text{tr}(P_N L P_N) + o(N^{1+2/n}),$$

where P_N denotes the spectral projection onto the first N eigenvalues.

(ii) **Semiclassical Weyl law.** Applying the semiclassical Weyl law [20, 15] to the symmetrized operator $L_s = (L + L^*)/2$ yields the leading-order term

$$\text{tr}(P_N L_s P_N) \sim C_n |\Omega| N^{1+2/n}.$$

Standard asymptotic techniques for such spectral sums can be found in [2].

(iii) **Control of the antisymmetric part.** The antisymmetric part $L_a = (L - L^*)/2$ contributes only at lower order, thanks to the growth bound in Assumption 2.4. Combining (i)–(iii) gives (5). $\square$

Corollary 3.2. *Under the hypotheses of Theorem 3.1, the counting function $N(\lambda) = \#\{k : \operatorname{Re} \lambda_k \leq \lambda\}$ satisfies*

$$N(\lambda) \sim \frac{\omega_n}{(2\pi)^n} |\Omega| \lambda^{n/2}, \quad \lambda \rightarrow \infty,$$

where ω_n is the volume of the unit ball in $\mathbb{R}^n$.

3.2 Pseudospectral bounds

Proposition 3.3. *Let L satisfy the hypotheses of Theorem 3.1, and let $\delta = \operatorname{dep}(L)$ be its departure from normality. Then*

$$\sigma_\varepsilon(L) \subseteq \sigma(L) + B(0, \varepsilon + C\delta^{1/2}), \quad (6)$$

where $B(0, r)$ denotes the disk of radius r centered at the origin.

Proof. Follows from [18, Theorem 4.3] combined with the growth estimates established in the proof of Theorem 3.1 and the perturbation bounds of. $\square$

Conjecture 3.4. *The bound (6) is sharp in the sense that the exponent $1/2$ cannot be improved to any smaller power.*

Problem 3.5 (Open). Extend Theorems 3.1 and 3.3 to the case of unbounded domains $\Omega = \mathbb{R}^n$, where the spectrum may contain continuous components.

■ 4 Numerical experiments

4.1 Setup

We consider the two-species system of [16]

$$\partial_t u = d_1 \Delta u + u(1 - u - av), \quad (7)$$

$$\partial_t v = d_2 \Delta v + v(bu - cv), \quad (8)$$

on the unit square $\Omega = (0, 1)^2$ with homogeneous Neumann boundary conditions. We use the parameters $a = 0.8$, $b = 1.2$, $c = 0.5$, $d_1 = 1$, and vary $d_2 \in [0.01, 1]$.

The linearized operator is discretized using a Chebyshev spectral method [3] with $N = 200$ collocation points in each direction, yielding a $2N^2 \times 2N^2$ matrix. The resulting generalized eigenvalue problem is solved by projecting onto the Grassmann manifold of k -dimensional subspaces, following the algorithms of [1].

4.2 Results

Figure 1 shows the spectrum and pseudospectrum contours for $d_2 = 0.05$. As predicted by Proposition 3.3, the pseudospectrum extends significantly beyond the spectrum, with the deviation growing as d_2 decreases. This behavior is consistent with the non-normality analysis of [8].




Figure 1. Spectrum (dots) and pseudospectrum contours ($\varepsilon = 10^{-1}, 10^{-2}, 10^{-3}$) of the linearized operator for $d_2 = 0.05$. The gold curve delimits the numerical range.

Table 1 compares the numerical eigenvalue sums with the asymptotic prediction (5). The agreement is excellent, with relative errors below 2% for $N \geq 50$.

N	Numerical $\sum \operatorname{Re} \lambda_k$	Asymptotic	Relative error
25	1.847×10^3	1.812×10^3	1.93%
50	7.402×10^3	7.285×10^3	1.61%
100	2.961×10^4	2.924×10^4	1.26%
200	1.184×10^5	1.172×10^5	1.02%

Table 1. Comparison of numerical eigenvalue sums with the asymptotic law (5).

Example 4.1. For the scalar case $m = 1$ and $d_1 = d_2$, the operator L is self-adjoint and (5) reduces to the classical Weyl law [20] with $C_n = (2\pi)^{-n} \omega_n$.

■ 5 Conclusions

We have established sharp spectral asymptotics for a class of non-self-adjoint reaction–diffusion operators. Our main results extend the classical Weyl law [20] to operators with controlled departure from normality and provide quantitative bounds on the pseudospectrum, in the spirit of [18].

Several directions remain open:

- (i) Extending the analysis to unbounded domains (Problem 3.5).
- (ii) Establishing sharpness of the pseudospectral bound (Conjecture 3.4).
- (iii) Applications to nonlinear pattern selection and Turing instability beyond the linear regime [19, 6].
- (iv) Connecting the spectral analysis to the structural biology of intrinsically disordered proteins, where non-normal effective operators arise naturally from conformational ensembles [5].

Acknowledgements

The authors thank the anonymous referees for valuable comments that improved the presentation of this work. This research was supported by Grant No. 12345 from the National Science Foundation and by the

RandPunkt Journal Early Career Award.

References

- [1] Absil, P.-A., Mahony, R., and Sepulchre, R., *Optimization Algorithms on Matrix Manifolds*, Princeton, NJ: Princeton University Press, 2008, doi:10.1515/9781400830244.
- [2] Bender, C. M. and Orszag, S. A., *Advanced Mathematical Methods for Scientists and Engineers*, Classical Theoretical Physics, New York: Springer-Verlag, 1999.
- [3] Boyd, J. P., *Chebyshev and Fourier Spectral Methods*, 2nd ed., Mineola, NY: Dover Publications, 2001.
- [4] Davies, E. B., *Spectral Theory and Differential Operators*, vol. 42, Cambridge Studies in Advanced Mathematics, Cambridge: Cambridge University Press, 1995, doi:10.1017/CB09780511623721.
- [5] Dunker, A. K. and Uversky, V. N., eds., *Intrinsically Disordered Proteins*, Hoboken, NJ: John Wiley & Sons, 2008.
- [6] Edelstein-Keshet, L., *Mathematical Models in Biology*, Classics in Applied Mathematics, Philadelphia: SIAM, 2005.
- [7] Evans, L. C., *Partial Differential Equations*, 2nd ed., vol. 19, Graduate Studies in Mathematics, Providence, RI: American Mathematical Society, 2010.
- [8] González, M. and Chen, L., *Pseudospectral Bounds for Advection–Diffusion Systems in Population Dynamics*, SIAM Journal on Applied Mathematics 84.3 (2024), pp. 1123–1147, doi:10.1137/23M1587241.
- [9] Hansen, P. C., *Rank-Deficient and Discrete Ill-Posed Problems: Numerical Aspects of Linear Inversion*, Monographs on Mathematical Modeling and Computation, Philadelphia: SIAM, 1998, doi:10.1137/1.9780898719697.
- [10] Hestenes, D. and Sobczyk, G., *Clifford Algebra to Geometric Calculus: A Unified Language for Mathematics and Physics*, Fundamental Theories of Physics, Dordrecht: D. Reidel Publishing Company, 1984, doi:10.1007/978-94-009-6292-7.
- [11] Keller, J. B., *Effective Behavior of Heterogeneous Media*, SIAM Review 19.2 (1977), pp. 261–285, doi:10.1137/1019042.
- [12] Lidskii, V. B., *On the Number of Solutions of the Equation $\det(\lambda I - A) = 0$* , Trudy Moskovskogo Matematicheskogo Obshchestva 8 (1959), pp. 199–222.
- [13] Maini, P. K., Painter, K. J., and Nguyen, H. P., *Spatial Pattern Formation in Chemical and Biological Systems*, Journal of the Chemical Society, Faraday Transactions 93.20 (1997), pp. 3601–3610, doi:10.1039/A702602A.
- [14] Murray, J. D., *Mathematical Biology II: Spatial Models and Biomedical Applications*, 3rd ed., New York: Springer-Verlag, 2003, doi:10.1007/b98869.
- [15] Reed, M. and Simon, B., *Methods of Modern Mathematical Physics IV: Analysis of Operators*, New York: Academic Press, 1978.
- [16] Segel, L. A. and Jackson, J. L., *Dissipative Structure: An Explanation and an Ecological Example*, Journal of Theoretical Biology 37.3 (1972), pp. 545–559, doi:10.1016/0022-5193(72)90090-2.
- [17] Smith, J. A. and Weber, K., *Semiclassical Weyl Laws for Non-Self-Adjoint Operators*, RandPunkt Journal of Applied Mathematics 1.1 (2023), pp. 1–28, doi:10.1000/rpjam.2023.001.
- [18] Trefethen, L. N. and Embree, M., *Spectra and Pseudospectra: The Behavior of Nonnormal Matrices and Operators*, Princeton, NJ: Princeton University Press, 2005.
- [19] Turing, A. M., *The Chemical Basis of Morphogenesis*, Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences 237.641 (1952), pp. 37–72, doi:10.1098/rstb.1952.0012.
- [20] Weyl, H., *Das asymptotische Verteilungsgesetz der Eigenwerte linearer partieller Differentialgleichungen*, Mathematische Annalen 71.4 (1911), pp. 441–479, doi:10.1007/BF01456804.